

CrIS observations of atmospheric composition in fire plumes over the Western US in summertime


<u>Vivienne Payne¹</u>, Susan Kulawik², Emily Fischer³, Kevin Bowman¹, Gene Francis⁴, Helen Worden⁴, Karen Cady-Pereira⁵, Frank Flocke⁴, Jakob Lindaas³, Ilana Pollack³


1. Jet Propulsion Laboratory, California Institute of Technology, 2. BAERI/NASA Ames, 3. Colorado State University, 4. National Center for Atmospheric Research, 5. Atmospheric and Environmental Research,

Acknowledgements:


JPL: Dejian Fu, Ming Luo, MUSES algorithm team
NASA Sounder Science Team
Sentinel Hub website

AIRS Science Team Meeting, Pasadena, April 2019










TES PAN retrievals are now available for the full TES record as part of the TES V7 Level 2 release.

TES and CrIS coverage



TES coverage for 8th November 2017

SNPP-CrIS coverage for 8th November 2017








Image credit: https://www.boiseweekly.com/

WE-CAN aircraft campaign: July 22nd to September 14th 2018

VIIRS image (NASA Worldview) S-NPP C112 -108 -120 -116 NOAA-20 E112 -120 -116 -108

740.pptv

580.pptv

420.pptv

260.pptv

100.pptv

740.pptv


580.pptv

420.pptv

260.pptv

100.pptv

Example WE-CAN flight day: 6th August 2018



Points show aircraft PAN measurments

Red line: 300 pptv

Western US is relatively cloud-free on this and other WE-CAN flight days. (Eastern area of this box is cloudy for this day.)

MUSES single footprint retrievals

Optimal estimation retrieval algorithm

- MUSES (Multi-species, multi-spectrum, multi-satellite)
- Single footprint (FOV) retrievals from L1B radiances

Forward model

- OSS (Optimal Spectral Sampling) from AER
- Trained on LBLRTM v12.4 with tes_v_3.0 line file


Sequential retrieval strategy


- PAN retrieval step follows other steps to fit for surface temperature, surface emissivity, cloud parameters, atmospheric temperature, H₂O, CO₂, O3
- MUSES also allows retrievals of other trace gases (e.g. CO)

Vertical sensitivity


- CrIS (and TES) sensitive mainly to PAN variations in the mid-upper troposphere
- Can show sensitivity to lower troposphere under certain conditions
- DOFS ~1.0

Vertical sensitivity: Example fire plume case







VIIRS image (NASA Worldview)




TROPOMI: Flying in constellation with S-NPP




TROPOMI aerosol index (Sentinel Hub website)
Thanks to Bryan Baum for pointing out existence of this product!




VIIRS image (NASA Worldview)




PAN DOF 20180806





VIIRS image (NASA Worldview)






20180806: PAN and CO


86.


180.pptv


100.pptv




TROPOMI aerosol index (Sentinel Hub website)

















Summary and next steps


- CrIS PAN retrievals for WE-CAN time period (Summer 2018)
 - 17 days processed with MUSES
 - High PAN values in fire plumes
 - Results look promising!
- Next steps:
 - Quantitative comparisons with WE-CAN aircraft "profiles"
 - WE-CAN flights extend up to ~6 km (~500 hPa), limited number of intentional profiles
 - Afternoon flights provide close match in time with satellite overpasses
 - Strongly elevated PAN values in smoke plumes
 - Quantitative comparisons with ATom-1 (Summer 2016) and ATom-2 (Spring 2017)
 - Aircraft profile measurements extend to ~13 km
 - Mostly "background" PAN values over remote ocean, some plumes observed
 - Temporal coincidences not necessarily ideal
 - Summer 2019: FIREChem and FIREX


Back-up slides

PAN shows spatial variation on small scales

Tight coincidences needed for validation

Example from FRAPPE campaign (20140729)





Modified from Fischer et al., 2018. ACP

https://www.star.nesdis.noaa.gov/JPSS/EDRs/products_Soundings_2018.php

